Subject index

Adirondack Mountains, 697
albite, 13
algebra, 251
Alleghanian, 145
aluminous reaction textures, 231
amphibole stability, 425
amphiboles, 539
amphibolite facies, 551
anchimetamorphism, 31
anchizone, 333
andalusite, 159, 199, 357, 375
anorthite = grossular + sillimanite, 493
Antarctica, 591
$^{40}Ar/^{39}Ar$ geochronology, 145
Archaean, 171, 591
Arunta Block, 65
Avalon Zone, 145

Bamble sector, south Norway, 231
barometry, 171, 683
basic schist, 425
biotite + sillimanite = garnet + K-feldspar, 493
blueschist facies, 345
brittle and ductile deformation, 457

Caledonides, 289
Capricorn Orogen, 47
central Australia, 65
central Shikoku, Japan, 413
clasts, 467
'clockwise' *P–T* path, 551
composition files, 251
conglomerate, 467
continuum mechanics, 13
coronas, 697
corundum + quartz, 125
crenulation cleavage, 13
Cr-rich staurolite, 231
crustal thickening, 257
crystal growth, 683
'crystallinity', 31

decompression, 591
deformation partitioning, 13
detachment faulting, 615
differential element mobility, 645
differential stress, 457
diffraction peak broadening, 31
ductile faults, 645

Eastern Alps, 629
eclogite, 289
epidote, 539
error and precision, 333
extensional models, 257

Fe^{2+}–Mg exchange (garnet–biotite), 493
Fiskefjord, southern West Greenland, 663
fluid chemistry, 131
fluid composition, 525
fluid inclusions, 131
fluid recirculation, 311
French Massif Central, 477
frictional melting, 507

garnet, 13, 357, 375, 683
garnet amphibolite, 467
garnet lherzolites, 477
garnet zoning, 551
geobarometry, 217
geological structure, 441
geothermobarometry, 269, 289, 525, 629
geothermometry, 217, 231
granite, 1
granulite facies, 551, 663
granulite facies metamorphism, 65
granulite metamorphism, 171
granulitic paragneisses, 477
Greenland, 171
greenschist facies, 345
Grenville gneisses, 493

hematite-bearing schist, 425
hercynitic spinel, 65
Hidaka metamorphic belt, 507
high fluid/rock ratios, 311
high temperature, 591
high thermal gradient, 159
high-*P*/*T* metamorphism, 401
high-pressure granulite, 289
high-temperature metamorphism, 551
hydrous minerals, 507

illite, 31
illite 'crystallinity', 333
intergranular diffusion, 697
internally consistent thermodynamic data set, 89
inverted metamorphism, 441
isocryst method, 333
isothermal decompression, 551

K-Ar, 289
KFMASH, 65
KFMASHTO, 65
Köli nappes, 615
kyanite, 231

Lachlan Fold Belt, 1
lack of porphyroblast rotation, 13
lithospheric thinning, 257
low-grade metamorphism, 131
low-*P* facies metamorphism, 257
low-pressure metamorphism, 1
low-temperature metamorphism, 539

Maine, 199
manganiferous iron formations, 525
Mary Kathleen Fold Belt, 311
metamorphic fluid flow, 311
metamorphic fluids, 131
metamorphic history, 269
metamorphic reactions, 697
metamorphic zonation, 413
metapelite, 289
metapelites, 357, 375
metasomatism, 645, 663
metastable equilibrium, 125
microcomputer programs, 251
microfossil, 401
microstructures, 629
migmatite, 1
migmatites, 551
mineral assemblages of metasediments, 217
multi-sample mass balance, 645

Namaqua Province, 159
Napier Complex, Antarctica, 125, 591

nappe, 441
Narryer Gneiss Complex, 47
'nested' *P–T–t* paths, 269
New England, 145
Newfoundland, Canada, 269
Northern Apennines, Ligurian Alps, 539
northern Australia, 311

orthoamphibole-zoning, 231
orthopyroxene–sillimanite, 591
Otago schists, 13
overstacking, 629
overstepping reactions, 125
oxidation, 375

pelitic schists, 683
petrogenetic grid, 375, 591
phase equilibrium calculations, 89
polymetamorphism, 289
projections, 251
pseudotachylites, 507
P–T estimate of late Alpine metamorphism, 217
P–T path, 47, 65, 231, 591
P–T paths, 615
P–T–D–X path, 375
P–T–t–d path, 629
P–T–t paths, 269, 289, 477
pumpellyite, 539
Puolankajärvi Formation, Finland, 357, 375
pyroxenes, 525

quench microlites, 507

radiometric age, 401
Rayner Complex, 591
Rb–Sr, 289
reaction history, 551
reactions, 251
reaction textures, 591
regional granulites, 47
regional metamorphism, 145, 257
retrograde metamorphism, 1
retrogression, 663
reworking, 591

Sambagawa, 457
Sambagawa belt, 413, 441
Sanbagawa schist, 425
Sanbagawa uplift, 467
sapphirine, 591
Scandinavian Caledonides, 615
sense of shear, 13
sequential growth, 13
sequential porphyroblast growth, 199
Shimanto, 457
sillimanite, 357, 375
Sm–Nd, 289
stable isotopes, 131
staurolite, 159, 199, 357, 375
structurally enhanced permeability, 311
structures, 357
Sudetes, 345
surface erosion, 401
syndeformational metamorphism, 199
syntectonic porphyroblasts, 159

Tauern Window, 217
terrane accretion, 145
textures, 357
thermal history, 145
thermal readjustment, 413
thermal structure, 413, 441
thermobarometry, 477
thermodynamic data, 89
thermometry, 683

uplift-cooling, 401
Uppermost Allochthon, 289

Variscan orogeny, 629
Variscides, 345

X-ray diffraction, 31

Yilgarn Block, 47

zoned garnets, 683

Author index

Andresen, A., *see* Krogh, E. J.
Andrews, M. S., *see* Dipple, G. M.

Ball, D. G. A. & Robin, P.-Y. F. METPET: Metamorphic petrology microcomputer programs, 251.
Banno, S., *see* Otsuki, M.
Banno, S., *see* Wallis, S.
Bevins, R. E., *see* Robinson, D.
Bhattacharya, A., Spiering, B., Sen, S. K., Natarajan, R. & Mazumdar, A. C. Compositional characteristics and phase equilibria in manganiferous iron formations from a high-grade terrain near Satnuru, Karnataka, India, 525.
Bhattacharya, P. K., *see* Sengupta, P.
Bottrell, S. H., Greenwood, P. B., Yardley, B. W. D., Shepherd, T. J. & Spiro, B. Metamorphic and post-metamorphic fluid flow in the low-grade rocks of the Harlech Dome, north Wales, 131.
Broks, T. M., *see* Krogh, E. J.
Brown, M., *see* Jones, K. A.
Bryhni, I., *see* Krogh, E. J.

Cabella, R., *see* Lucchetti, G.
Carlson, W. D., *see* Johnson, C. D.
Clarke, G. L., Collins, W. J. & Vernon, R. H. Successive overprinting granulite facies metamorphic events in the Anmatjira Range, central Australia, 65.
Collins, W. J., *see* Clarke, G. L.
Cortesogno, L., *see* Lucchetti, G.
Crowley, P. D. Metamorphism within the Çokkul synform: evidence for detachment faulting within the metamorphic infrastructure of the Norwegian Caledonides (67°30′N), 615.

Dachs, E. Geothermobarometry in metasediments of the southern Grossvenediger area (Tauern Window, Austria), 217.
Dallmeyer, R. D., Hermes, O. D. & Gil-Ibarguchi, J. I. $^{40}Ar/^{39}Ar$ mineral ages from the Scituate Granite, Rhode Island: implications for Late Palaeozoic tectonothermal activity in New England, 145.
Dasgupta, S., *see* Sengupta, P.
Dipple, G. M., Wintsch, R. P. & Andrews, M. S. Identification of the scales of differential element mobility in a ductile fault zone, 645.
Dufour, E., *see* Gardien, V.
Dunn, G. R., *see* Lang, H. M.

Etheridge, M. A., *see* Loosveld, R. J. H.

Florence, F. P., *see* Spear, F. S.

Garde, A. A. Thermal granulite facies metamorphism with diffuse retrogression in Archaean orthogneisses, Fiskefjord, southern West Greenland, 663.
Gardien, V., Tegyey, M., Lardeaux, J. M., Misseri, M. & Dufour, E. Crust–mantle relationships in the French Variscan chain: the example of the Southern Monts du Lyonnais unit (eastern French Massif Central), 477.
Gil-Ibarguchi, J. I., *see* Dallmeyer, R. D.
Greenwood, P. B., *see* Bottrell, S. H.

Hara, I., Shiota, T., Hide, K., Okamoto, K., Takeda, K., Hayasaka, Y. & Sakurai, Y. Nappe structure of the Sambagawa belt, 441.
Harley, S. L., Hensen, B. J. & Sheraton, J. W. Two-stage decompression in orthopyroxene–sillimanite granulites from Forefinger Point, Enderby Land, Antarctica: implications for the evolution of the Archaean Napier Complex, 591.
Hayasaka, Y., *see* Hara, I.
Hensen, B. J., *see* Motoyoshi, Y.
Hensen, B. J., *see* Harley, S. L.
Hermes, O. D., *see* Dallmeyer, R. D.
Hide, K., *see* Hara, I.
Higashino, T. The higher grade metamorphic zonation of the Sambagawa metamorphic belt in central Shikoku, Japan, 413.
Holland, T. J. B. & Powell, R. An enlarged and updated internally consistent thermodynamic dataset with uncertainties and correlations: the system K_2O–Na_2O–CaO–MgO–MnO–FeO–Fe_2O_3–Al_2O_3–TiO_2–SiO_2–C–H_2–O_2, 89.
Humphreys, H. C. & Van Bever Donker, J. M. Early Namaqua low-pressure metamorphism: deformation and porphyroblast growth in the Zoovoorby staurolite schist, South Africa, 159.

Isozaki, Y. & Itaya, T. Chronology of Sanbagawa metamorphism, 401.
Itaya, T., *see* Isozaki, Y.
Itaya, T., *see* Yokoyama, K.

Jamieson, R. A. Metamorphism of an Early Palaeozoic continental margin, western Baie Verte Peninsula, Newfoundland, 269.
Johnson, C. D. & Carlson, W. D. The origin of olivine–plagioclase coronas in metagabbros from the Adirondack Mountains, New York, 697.
Johnson, S. E. Lack of porphyroblast rotation in the Otago schists, New Zealand: implications for crenulation cleavage development, folding and deformation partitioning, 13.
Jones, K. A. & Brown, M. High-temperature 'clockwise' *P–T* paths and melting in the development of regional migmatites: an example from southern Brittany, France, 551.

Kisch, H. J. Calibration of the anchizone: a critical comparison of illite 'crystallinity' scales used for definition, 31.
Kohn, M. J., *see* Spear, F. S.
Kretz, R. Biotite and garnet compositional variation and mineral equilibria in Grenville gneisses of the Otter Lake area, Quebec, 493.
Kristensen, S. E., *see* Krogh, E. J.
Krogh, E. J., Andresen, A., Bryhni, I., Broks, T. M. & Kristensen, S. E. Eclogites and polyphase *P–T* cycling in the Caledonian Uppermost Allochthon in Troms, northern Norway, 289.
Kryza, R., Muszynski, A. & Vielzeuf, D. Glaucophane-bearing assemblage overprinted by greenschist-facies metamorphism in the Variscan Kaczawa complex, Sudetes, Poland, 345.

Laajoki, K. & Tuisku, P. Metamorphic and structural evolution of the Early Proterozoic Puolankajärvi Formation, Finland – I. Structural and textural relations, 357.
Laajoki, K., *see* Tuisku, P.
Lang, H. M. & Dunn, G. R. Sequential porphyroblast growth during deformation in a low-pressure metamorphic terrain, Orrs Island–Harpswell Neck, Maine, 199.
Lardeaux, J. M., *see* Gardien, V.
Loosveld, R. J. H. & Etheridge, M. A. A model for low-pressure facies metamorphism during crustal thickening, 257.
Lucchetti, G., Cabella, R. & Cortesogno, L. Pumpellyites and coexisting minerals in different low-grade metamorphic facies of Liguria, Italy, 539.

Matsueda, H., *see* Motoyoshi, Y.
Mazumdar, A. C., *see* Bhattacharya, A.
McGregor, V. R., *see* Riciputi, L. R.
Menard, T., *see* Spear, F. S.
Misseri, M., *see* Gardien, V.
Morand, V. J. Low-pressure regional metamorphism in the Omeo Metamorphic Complex, Victoria, Australia, 1.
Motoyoshi, Y., Hensen, B. J. & Matsueda, H. Metastable growth

of corundum adjacent to quartz in a spinel-bearing quartzite from the Archaean Napier Complex, Antarctica, 125.
Muhling, J. R. The Narryer Gneiss Complex of the Yilgarn Block, Western Australia: a segment of Archaean lower crust uplifted during Proterozoic orogeny, 47.
Mukherjee, M., *see* Sengupta, P.
Muszynski, A., *see* Kryza, R.

Natarajan, R., *see* Bhattacharya, A.

Okamoto, K., *see* Hara, I.
Oliver, N. H. S., Valenta, R. K. & Wall, V. J. The effect of heterogeneous stress and strain on metamorphic fluid flow, Mary Kathleen, Australia, and a model for large-scale fluid circulation, 311.
Otsuki, M. & Banno, S. Prograde and retrograde metamorphism of hematite-bearing basic schists in the Sanbagawa belt in central Shikoku, 425.

Powell, R., *see* Holland, T. J. B.

Riciputi, L. R., Valley, J. W. & McGregor, V. R. Conditions of Archean granulite metamorphism in the Godthab–Fiskenaesset region, southern West Greenland, 171.
Robin, P.-Y. F., *see* Ball, D. G. A.
Robinson, D., Warr, L. N. & Bevins, R. E. The illite 'crystallinity' technique: a critical appraisal of its precision, 333.

Sakurai, Y., *see* Hara, I.
Schulz, B. Prograde–retrograde P–T–t–deformation path of Austroalpine micaschists during Variscan continental collision (Eastern Alps), 629.
Sen, S. K., *see* Bhattacharya, A.
Sengupta, P., Dasgupta, S., Bhattacharya, P. K. & Mukherjee, M. An orthopyroxene–biotite geothermometer and its application in crustal granulites and mantle-derived rocks, 191.
Senior, A., *see* Visser, D.
Shepherd, T. J., *see* Bottrell, S. H.
Sheraton, J. W., *see* Harley, S. L.
Shiota, T., *see* Hara, I.
Spear, F. S., Kohn, M. J., Florence, F. P. & Menard, T. A model for garnet and plagioclase growth in pelitic schists: implications for thermobarometry and P–T path determinations, 683.
Spiering, B., *see* Bhattacharya, A.
Spiro, B., *see* Bottrell, S. H.

Takeda, K., *see* Hara, I.
Tegyey, M., *see* Gardien, V.
Toriumi, M. The transition from brittle to ductile deformation in the Sambagawa metamorphic belt, Japan, 457.
Toyoshima, T. Pseudotachylite from the Main Zone of the Hidaka metamorphic belt, Hokkaido, northern Japan, 507.
Tuisku, P. & Laajoki, K. Metamorphic and structural evolution of the Early Proterozoic Puolankajärvi Formation, Finland – II. The pressure–temperature–deformation–composition path, 375.
Tuisku, P., *see* Laajoki, K.

Valenta, R. K., *see* Oliver, N. H. S.
Valley, J. W., *see* Riciputi, L. R.
Van Bever Donker, J. M., *see* Humphreys, H. C.
Vernon, R. H., *see* Clarke, G. L.
Vielzeuf, D., *see* Kryza, R.
Visser, D. & Senior, A. Aluminous reaction textures in orthoamphibole-bearing rocks: the pressure–temperature evolution of the high-grade Proterozoic of the Bamble sector, south Norway, 231.

Wall, V. J., *see* Oliver, N. H. S.
Wallis, S. & Banno, S. The Sambagawa belt – trends in research, 393.
Warr, L. N., *see* Robinson, D.
Wintsch, R. P., *see* Dipple, G. M.

Yardley, B. W. D., *see* Bottrell, S. H.
Yokoyama, K. & Itaya, T. Clasts of high-grade Sanbagawa schist in Middle Eocene conglomerates from the Kuma Group, central Shikoku, south-west Japan, 467.